

# NMR Analysis of Oligosaccharides Containing Fructopyranoside

Eri Fukushi<sup>1\*</sup> • Hideki Okada<sup>2</sup> • Akira Yamamori<sup>2</sup> • Naoki Kawazoe<sup>2</sup> •  
Shuichi Onodera<sup>3</sup> • Jun Kawabata<sup>1</sup> • Norio Shiomi<sup>3</sup>

<sup>1</sup> Graduate school of Agriculture, Hokkaido University, Sapporo 060-8589, Japan

<sup>2</sup> Ohtakakohso, Co. Ltd., Otaru 047-0193, Japan

<sup>3</sup> Department of Food and Nutrition Science, Graduate School of Dairy Science Research, Rakuno Gakuen University, Ebetsu 069-8501, Japan

Corresponding author: \* feria@cen.agr.hokudai.ac.jp

## ABSTRACT

This review focuses on the NMR methods for the oligosaccharides containing fructopyranoside that were previously isolated from the fermented beverage of an extract from 50 kinds of fruits and vegetables. The <sup>1</sup>H and <sup>13</sup>C-NMR signals of each saccharide were assigned using 2D-NMR including COSY, HSQC, HSQC-TOCSY, CH<sub>2</sub>-selected HSQC-TOCSY, and CT (constant time)-HMBC. The fructose in pyranosyl form showed different <sup>13</sup>C chemical shifts from those of furanosyl form. Further confirmation of the pyranosyl form could be obtained from the HMBC correlation peak between C-2 and H-6 of fructose residue (Fru), whereas the C-2 of Fru in furanosyl form could give the HMBC correlation peak between H-5 of Fru. Problems encountered were signal overlapping of protons and low peak separation. The key correlation peak between C-2 and H-6 of Fru was overlapped by the correlation peak indicating a glycosidic linkage between the C-2 of Fru and the H-6 of the glucose residue (Glc, or Glc'). These were solved using HSQC and CT-HMBC spectra rather than HMQC and conventional HMBC spectra, which have an inherent broad-line-shape in the carbon dimension.

**Keywords:** COSY, HSQC, HSQC-TOCSY, CH<sub>2</sub>-selected HSQC-TOCSY, CT-HMBC

**Abbreviations:** COSY, correlated spectroscopy; CT-HMBC, constant time-HMBC; E-HSQC, editing-HSQC; HMBC, heteronuclear multiple bond correlation; HSQC, heteronuclear single quantum coherence; TOCSY, total correlation spectroscopy

## CONTENTS

INTRODUCTION.....	10
General procedure for NMR analysis .....	10
NMR analysis with special methods for high resolution in the <sup>13</sup> C axis .....	13
PERSPECTIVES AND CONCLUSIONS.....	13
REFERENCES.....	13

## INTRODUCTION

The NMR spectra of oligosaccharides are too complex to analyze using routine methods, COSY, HSQC, HSQC-TOCSY, and HMBC. The problems are signal overlapping of protons and low peak separation. The key HMBC correlation peak, which indicates a sugar-linkage, tends to be overlapped by intra-residual correlation peaks. Therefore, we created special methods to overcome this problem. Several oligosaccharides containing fructopyranoside were isolated from fermented beverage of an extract from 50 kinds of fruits and vegetables (Okada *et al.* 2006; Kawazoe *et al.* 2008). Their structure had been analyzed in a same procedure by using 1D and 2D NMR methods. Since the conventional NMR methods could not complete the analysis of their structure, the high-resolution 2D methods were applied. The established protocol performed the structural analysis of related oligosaccharide. In this paper, the procedure of these NMR analysis and efficiency of utilization of advanced NMR methods will be demonstrated using saccharide **1** (Fig. 1) as a model compound.

### General procedure for NMR analysis

**Fig. 2A** shows a one-dimensional (1D) carbon (<sup>13</sup>C)-NMR spectrum of saccharide **1**. NMR detects the nuclear magnetic resonance while 1D NMR spectrum provides a record about one kind of nucleus, in this case, the nucleus of car-






Fig. 1 Structure of saccharide **1**.

bon. The horizontal axis, which indicates the chemical shift (in ppm) is an important value of NMR in which each carbon placed in a different environment, shows a different chemical shift. **Fig. 2B** is a 1D proton (<sup>1</sup>H)-NMR spectrum. The fine structure of the signal because of the proton-proton coupling indicates information about neighboring protons. The anomer configuration of  $\alpha$ - or  $\beta$ -aldose can be determined by the coupling constant (splitting width) of the anomeric proton.

One-dimensional <sup>1</sup>H and <sup>13</sup>C NMR signals are assigned using two-dimensional (2D) NMR spectroscopy. **Fig. 3** shows a 2D HSQC (Bodenhausen *et al.* 1980; Willker *et al.* 1993) spectrum. The first dimension ( $F_1$ ), the vertical axis, is a carbon chemical shift ( $\delta_C$ ). The second dimension ( $F_2$ ), the horizontal axis, is a proton chemical shift ( $\delta_H$ ). Proton and carbon 1D spectra are attached. The HSQC detects the



**Fig. 2** One dimensional (1D)  $^{13}\text{C}$ (A, 126 MHz)- and  $^1\text{H}$ (B, 500MHz)-NMR spectra of **1** (5 mg / 0.5 ml of  $\text{D}_2\text{O}$ ). Chemical shifts of  $^1\text{H}$  ( $\delta_{\text{H}}$ ) and  $^{13}\text{C}$  ( $\delta_{\text{C}}$ ) in ppm were determined relative to the external standard of sodium [2,2,3,3- $^2\text{H}_4$ ]-3-(trimethylsilyl) propanoate in  $\text{D}_2\text{O}$  ( $\delta_{\text{H}}$  0.00 ppm) and 1,4-dioxane ( $\delta_{\text{C}}$  67.40 ppm) in  $\text{D}_2\text{O}$ .



**Fig. 3** Two dimensional (2D) HSQC spectrum of **1**.




Fig. 4 2D HSQC-TOCSY spectrum of **1**.

directly bonded proton and carbon. For example, the cross peak marked by an arrow is on the crossing point of the horizontal line from the carbon at  $\delta_C$  92.71 ppm (C-1 of  $\alpha$ Glc) and the vertical line from the proton at  $\delta_H$  at 5.25 ppm (H-1 of  $\alpha$ Glc). This means that this proton and this carbon are connected directly. The NMR spectra of saccharide **1** show that it was an anomeric mixture at the glucose residue. The  $\beta$  anomer is predominant and some signals of the  $\alpha$  anomer were distinct and could be assigned.

Fig. 4 shows a 2D  $^1\text{H}$ - $^{13}\text{C}$  HSQC-TOCSY (Willker *et al.* 1993; Domke *et al.* 1991). It is the combined method of HSQC and TOCSY (Braunschweiler *et al.* 1983; Hurd 1990; Parella *et al.* 1997). Each protonated carbon gives a HSQC peak for the directly coupled proton, and in the same row at this carbon, additional peaks for protons belonging the same spin-spin network with that proton. The HSQC-TOCSY spectrum can assign protons and carbons in the same spin-system. In an oligosaccharide spectrum, almost every signal in the same sugar unit can be identified. This spectrum revealed the  $^1\text{H}$ -spin systems of each sugar residue. The isolated methylene was assigned as H-1 and C-1 of Fru. The quaternary carbon was assigned as C-2 of Fru.

Fig. 5 is the 2D COSY (Aue *et al.* 1975; von Kienlin *et al.* 1991) spectrum. It detects protons via two or three bonds. There is a scalar coupling ( $J$  coupling) between protons via two or three bonds. The NMR parameters,  $J$  values (in Hz) of such scalar couplings depend on the stereochemistry, can be used to determine the sugar type and its anomeric configuration. The starting signals to analyze this COSY spectrum are the anomeric protons (H-1) of glucoses. They are characteristic doublet signals at rather large chemical shifts in the  $^1\text{H}$ -NMR spectrum. This spectrum is assigned from H-1 to H-3 in each glucose residue. Other signals could not be assigned because of signal overlapping.

2D  $^1\text{H}$ - $^{13}\text{C}$  HMBC (Bax *et al.* 1986; Summers *et al.* 1986; Hurd *et al.* 1991; Furihata *et al.* 1998) detects long-range  $J_{\text{CH}}$  couplings. This can also detect the connectivity between proton and carbon via oxygen and quaternary carbon. It is a useful tool in determining the glycosidic linkages and assigning the fructose unit. In the Fru unit, the HMBC correlations between C-3 and H-1, and C-2 and H-1




Fig. 5 2D COSY spectrum of **1**.

confirm the assignment of these signals (Fig. 6). The other four methylene carbons were assigned as C-6 in these residues. As for glucose residues, the HMBC correlations between C-5 and H-1 revealed the assignment of C-5 in each residue. The remaining  $^1\text{H}$  and  $^{13}\text{C}$  signals in each glucose residues could be assigned as H-4 and C-4. These results clarified the assignment of  $^1\text{H}$  and  $^{13}\text{C}$ -NMR signals of each residue.

The position of the glycosidic linkage, and pyranoside-form of Fru were analyzed as follows. The C-3 in  $\beta$ Glc and  $\alpha$ Glc showed HMBC correlations between H-1 in Glc', and C-1 in Glc' showed the HMBC correlations to H-3 in  $\beta$ -Glc and  $\alpha$ -Glc (Fig. 6). The  $J$ -value taken from the 1D  $^1\text{H}$ -NMR spectrum between H-1 and H-2 in Glc' was 7.7 Hz. These results indicate **1** has the Glc'  $\alpha$ 1- $\rightarrow$ 3 Glc linkage, namely, the laminaribiose moiety. The C-2 of Fru showed the




Fig. 6 2D HMBC spectrum of **1**.

HMBC correlations to H-6 of Glc' and H-6 of Fru (Fig. 6). These results revealed the fructopyranoside residue and its 2->6 Glc' linkage. The anomeric configuration and pyranosyl/furanosyl forms can be confirmed by the  $^{13}\text{C}$  chemical shifts (Shiomi *et al.* 2007).

### NMR analysis with special methods for high resolution in the $^{13}\text{C}$ axis

The major problem of NMR of oligosaccharides is signal overlapping. In general,  $^1\text{H}$  and  $^{13}\text{C}$ -NMR signals spread at 0–10 and 0–220 ppm, respectively. However, in typical saccharides, they are concentrated at 3–5 and 60–110 ppm, respectively. Especially methylene carbons are concentrated in a few ppm. For determination of sugar linkages and fructopyranoside form, the assignment of each methylene was necessary. So, we created special methods for higher resolution in the  $^{13}\text{C}$  dimension. That is,  $\text{CH}_2$ -selected (Fukushi *et al.* 2000) Editing(E)-HSQC (Willker *et al.* 1993; Davis *et al.* 1991) and  $\text{CH}_2$ -selected E-HSQC-TOCSY (Okada *et al.* 2003).

Peak separation is dependent on the spectral resolution. This is one of the key factors of spectral quality. For example, the expansion plot of the indicated region of the Fig. 7A is shown in Fig. 7B. In B, the correlation peaks also became larger so these peaks continue to overlap. The resolution is defined as spectral width per data point: the smaller the value, the higher the resolution. An increase of data points in the  $^{13}\text{C}$  axis is effective for improving the resolution of the  $^{13}\text{C}$  dimension. Narrowing the spectral width of the  $^{13}\text{C}$  dimension is also effective for improving the resolution. However, the spectral width of the  $^{13}\text{C}$  dimension for a conventional HSQC spectrum is usually set to cover the range of all protonated carbons. If there are signals outside the measuring range, they appear in the

spectrum as unwanted folded signals.

The E-HSQC gives each  $\text{CH}$ -,  $\text{CH}_2$ -, and  $\text{CH}_3$ -subpectrum. The spectral width of the  $^{13}\text{C}$  dimension and data points can be optimized for each subspectrum. The chemical shifts of methylene carbons for compound **1** are limited to a narrow region, enabling the spectral range of the  $^{13}\text{C}$  axis to be set narrower, so that a high resolution results. The C-6 in each residue can be assigned by this  $\text{CH}_2$ -selected E-HSQC spectrum (Fig. 8). In each glucose residue, the correlation peak appeared separately between C-6 and H-1.

### PERSPECTIVES AND CONCLUSIONS

The NMR methods for the oligosaccharides containing fructopyranoside are demonstrated above. The NMR spectra of oligosaccharides are too complex to analyze using routine methods, COSY, HSQC, HSQC-TOCSY, and HMBC. Problems of signal overlapping and low peak separation can be solved using high resolution  $\text{CH}_2$ -selected HSQC-TOCSY, and CT (constant time)-HMBC spectra. The procedure of these NMR analyses is also efficient for other kinds of compounds.

### REFERENCES

- Aue WP, Batholdi E, Ernst RR (1975) Two-dimensional spectroscopy. Application to nuclear magnetic resonance. *Journal of Chemical Physics* **64**, 2229-2246
- Bax A, Summers MF (1986)  $^1\text{H}$  and  $^{13}\text{C}$  assignments from sensitivity-enhanced detection of heteronuclear multiple-bond connectivity by 2D multiple quantum NMR. *Journal of the American Chemical Society* **108**, 2093-2094
- Bodenhausen G, Ruben DJ (1980) Natural abundance nitrogen-15 NMR by enhanced heteronuclear spectroscopy. *Chemical Physics Letters* **69**, 185-189
- Braunschweiler L, Ernst RR (1983) Coherence transfer by isotropic mixing: application to proton correlation spectroscopy. *Journal of Magnetic Resonance* **53**, 521-528




Fig. 7 2D HSQC-TOCSY (A) of 1 and expansion plot (B) of indicated area of (A).




Fig. 8  $\text{CH}_2$ -selected E-HSQC-TOCSY of 1.

Davis DG (1991) Improved multiplet editing of proton-detected, heteronuclear shift-correlation spectra. *Journal of Magnetic Resonance* **91**, 665-672

Domke T (1991) A new method to distinguish between direct and remote signals in proton-relayed X, H correlations. *Journal of Magnetic Resonance* **95**, 174-177

Fukushi E, Onodera S, Yamamori A, Shiomi N, Kawabata J (2000) NMR analysis of tri- and tetrasaccharides from asparagus. *Magnetic Resonance in Chemistry* **38**, 1005-1011

Furihata K, Seto H (1998) Constant time HMBC (CT-HMBC), a new HMBC technique useful for improving separation of cross peaks. *Tetrahedron Letters*

**39**, 7337-7340

Hurd RE (1990) Gradient-enhanced spectroscopy. *Journal of Magnetic Resonance* **87**, 422-428

Hurd RE, John BK (1991) Gradient-enhanced proton-detected heteronuclear multiple-quantum coherence spectroscopy. *Journal of Magnetic Resonance* **91**, 648-653

Kawazoe N, Okada H, Fukushi E, Yamamori A, Onodera S, Kawabata J, Shiomi N (2008) Two novel oligosaccharides isolated from a beverage produced by fermentation of a plant extract. *Carbohydrate Research* **343**, 549-554

- Okada H, Fukushi E, Onodera S, Nishimoto T, Kawabata J, Kikuchi M, Shiomi N** (2003) Synthesis and structural analysis of five novel oligosaccharides prepared by glucosyl transfer from  $\beta$ -D-glucose 1-phosphate to isokestose and nystose using *Thermoanaerobacter brockii* kojibiose phosphorylase. *Carbohydrate Research* **338**, 879-885
- Okada H, Fukushi E, Yamamori A, Kawazoe N, Onodera S, Kawabata J, Shiomi N** (2006) Structural analysis of a novel saccharide isolated from fermented beverage of plant extract. *Carbohydrate Research* **341**, 925-929
- Parella T, Sánchez-Ferrando F, Virgili A** (1997) Quick recording of pure absorption 2D TOCSY, ROESY, and NOESY spectra using pulsed field gradients. *Journal of Magnetic Resonance* **125**, 145-148
- Shiomi N, Benkeblia N, Onodera S** (2007) *Recent Advances in Fructooligosaccharides Research*, Research Signpost, Kerala, India, pp 119-134
- Summers MF, Marzilli LG, Bax A** (1986) Complete  $^1\text{H}$  and  $^{13}\text{C}$  assignments of coenzyme B<sub>12</sub> through the use of new two-dimensional NMR experiments. *Journal of the American Chemical Society* **108**, 4285-4294
- von Kienlin M, Moonen CTW, van der Toorn A, van Zijl PCM** (1991) Rapid recording of solvent-suppressed 2D COSY spectra with inherent quadrature detection using pulsed field gradients. *Journal of Magnetic Resonance* **93**, 423-429
- Willker W, Leibfritz D, Kerssebaum R, Bermel W** (1993) Gradient selection in inverse heteronuclear correlation spectroscopy. *Magnetic Resonance in Chemistry* **31**, 287-292